

DISCHARGE INSTRUCTIONS AFTER PORTACATH PLACEMENT

WHAT TO EXPECT AFTER SURGERY

- You should expect a fair amount of pain at the incision site. This should improve quickly.
- You will feel a lump under the skin- it is the port. This is normal.

DIET AND BOWEL HABITS: Eat your usual diet with plenty of fruits and vegetables and bran cereal. Drink about 8 glasses of fluid/water daily. We strongly recommend a fiber supplement like Citrucel or Metamucil taken once or twice daily. You preferably should have a soft, yet formed BM daily without straining. You should take a laxative while on narcotics. Options include Miralax 1 or more tablespoons daily, Kondremul 1-2 tablespoons daily or Dulcolax 2 tablespoons 2 times per day until you have a BM. If you are having liquid BM's, then stop.

CARE OF SURROUNDING SKIN: You may wash with soap and water. Please watch for signs of infection including redness, swelling and drainage. Call the office for further instructions if this occurs.

ACTIVITY: This is determined by your comfort level. Driving should wait until at least 48 hours after surgery and no narcotics within 8 hours of driving. You should only do light activity for 2 weeks while the port secures into place.

FOLLOW-UP: Please call the office at **503-281-0561** to schedule a follow-up visit if you do not have any plans to see your Oncologist in the next 2 weeks. Otherwise, the Oncologist and his staff will monitor the port site. However, if you are experiencing increasing pain, fever (>101F), or any other concerning issues call to be seen sooner.

David O'Brien, MD, FACS, FASCRS

Colon and Rectal Surgery

The Oregon Clinic

4805 NE Glisan Street, Suite 6N60

Portland, OR 97213

(503) 281-0561

Instructed by: _____

Released To: _____ Date: _____